


EXERCISE SOLUTION

- 1(a) Yes, because the size of railway carriage is very small as compared to the distance between two stations.
- (b) Yes, because monkey is very small as compared to the circular track.
- (c) No, because turning of ball is not smooth. Thus the distance covered by the ball is not large enough to consider the ball as a point object.
- (d) No, because the distance covered by the tumbling beaker is not much larger.

2.



- (a) Because $OP < OQ$, hence A lives closer to school than B.
- (b) When $x=0$, $t=0$ for A but for B, t has some non-zero value. Therefore.

A starts from the school earlier than B.

(c) Slope of graph for B > slope for A. Therefore
Speed of B > Speed of A. i.e.
B walks faster than A

(d) Since 'it' is same for both A and B for P
and a position, we can see.
A and B reach home at the same time.

(e) x-t graph for A and B intersect only
once. Hence
B overtakes A on the road once.

3. Time taken by woman to reach the office

$$= \frac{\text{distance}}{\text{speed}} = \frac{s}{v}$$

$$= \frac{2.5}{5} = \frac{1}{2} \text{ hr} = 30 \text{ min}$$

Therefore she reaches the office at 9.30 a.m.
She stays at the office between 9.30 to 5.00 p.m.
Now


time taken for the return journey

$$= \frac{\text{distance}}{\text{speed}}$$

$$= \frac{2.5}{2.5} = \frac{1}{10} \text{ hr} = 6 \text{ min.}$$

Hence she reaches at home at 5.06 p.m

The $x-t$ graph for the woman's motion



4.

The drunkard takes 5 steps forward and 3 steps backward and so on.

Each step is 1 m long and requires time 1 sec.
So,

time taken to travel 2 m = 8 sec.


$\therefore$ time taken to travel 8 m = $\frac{8 \times 8}{2} = 32$ sec.

In the last 5 sec man will cover 5 steps. i.e.
the distance = 13 m (= 8 + 5) m

and time taken to cover 13 m = $32 + 5$
= 37 sec.

Hence man will take 37 sec to fall into a pit 13 m away.

The $x-t$ graph for the motion of drunkard



3.6

Initial speed of the car $u = 126 \text{ km/h}$

$$= \frac{126 \times 1000}{60 \times 60} \text{ m/s}$$

$$= 126 \times \frac{5}{18}$$

$$= 7 \times 5$$

$$u = 35 \text{ m/s}$$

Final speed $v = 0$ distance covered $s = 200 \text{ m}$

Use

$$v^2 - u^2 = 2as$$

$$(0)^2 - (35)^2 = 2 \times a \times 200$$

$$\text{or } a = \frac{-(35)^2}{2 \times 200} = \frac{-1225}{400}$$

$$= -3.06 \text{ m/s}^2 \quad \underline{\text{Ans}}$$

Now we use

$$v = u + at$$


$$0 = 35 - 3.06 \times t$$

$$\text{or } t = \frac{35}{3.06} = 11.4 \text{ sec}$$

Ans

3.9 x (not in new book)

NCERT Exercise



Let the speed of bus $= V_B$
 Distance travelled by bus in $T_{min} = L$
 Speed of cyclist $V_C = 20 \text{ km/h}$

For the buses leaving from town A

$$L = V_B \times \frac{T}{60} \text{ km} \quad [\because T_{min} = \frac{T}{60} \text{ hrs}]$$

w.r. to cyclist

$$L = (V_{Bc}) \times \frac{18}{60} \quad [V_{Bc} = \text{Relative speed of bus w.r. to cyclist}]$$

$$\text{or } L = (V_B - V_C) \times \frac{18}{60} \quad \text{--- (1)} \quad [V_B \text{ and } V_C \text{ have same direction}]$$

For the buses leaving from town B

$$L = V_B \times \frac{T}{60}$$

w.r. to cyclist

$$L = V_{Bc} \times \frac{6}{60}$$

$$L = (V_B + V_C) \times \frac{6}{60} \quad \text{--- (2)} \quad [V_B \text{ and } V_C \text{ have opposite direction}]$$

From eqⁿ (1) and (2)

$$(V_B - V_C) \frac{18}{60} = (V_B + V_C) \frac{6}{60}$$

$$\text{or } 3(V_B - V_C) = V_B + V_C$$

$$\text{or } 3V_B - V_B = 3V_C + V_C$$

$$\text{or } 2V_B = 4V_C$$

$$\text{or } V_B = 2V_C$$

$$= 2 \times 20 \quad [\because V_C = 20 \text{ km/h}]$$

$$V_B = 40 \text{ km/h} \quad \underline{\text{Ans}}$$

Now put the value of V_B in relation

$$L = V_B \times \frac{T}{60}$$

$$L = (V_B - V_C) \times \frac{18}{60} \quad [\text{eq}^n (1)]$$

$$L = (40 - 20) \times \frac{18}{60} = 20 \times \frac{18}{60} = 6$$

$$\text{or } L = 6 \text{ km}$$

$$\text{Now by using } L = V_B \times \frac{T}{60}$$

$$T = \frac{60 \times L}{V_B} = \frac{60 \times 6}{40}$$

$$T = 9 \text{ min}$$

Ans

2.6

Topic.....

Date.....

3.10

(a)

for vertical motion

Acceleration due to gravity (g) is always in downward direction.

(b)

Velocity at highest point

$$v=0$$

and $g = 9.8 \text{ m/s}^2$ vertically downwards

(c)

During upward motion

(i) Position is +ve

(ii) velocity is -ve

(iii) accⁿ due to gravity g is +ve ($a=g$)

During downward motion

(i) Position is +ve

(ii) Velocity is +ve

(iii) g is +ve(d) $u = -29.4 \text{ m/s}$, $v=0$, $a=g=9.8 \text{ m/s}^2$ by $v^2 = u^2 + 2a(x-x_0)$

$$0 = (-29.4)^2 + 2 \times 9.8 (x-x_0)$$




$$\text{or } x-x_0 = \frac{29.4 \times 29.4}{2 \times 9.8}$$

$$\text{or } x-x_0 = \frac{-0.82}{2} = -44.1$$

$$\text{or } 0-x_0 = -44.1 \quad [x=0]$$

$$\text{or } x_0 = 44.1 \text{ m}$$

i.e. ball rise upto 44.1 m.



Teacher's Sign

Also,

$$v = u + at$$

$$\therefore 0 = -29.4 + 9.8 \times t$$

$$\text{or } t = 3 \text{ sec}$$

total time in which ball return to the player's hand = $2t$

$$= 2 \times 3 = 6 \text{ sec} \quad \underline{Ans}$$

2.7

3.11

(a) True

At highest point of the motion the body is at rest for a moment but the acceleration due to gravity (g) is vertically downwards.

(b) False

If a particle has non-zero velocity, it must have speed.

(c) True

When a particle is moving with uniform velocity acceleration is zero. ($\because a = \frac{\Delta v}{\Delta t}$)

(d) False

It depends on the coordinate system taken to explain the motion of the particle.

2.8

Topic.....

Date.....

3.12

When the ball falls from
90 m height -
(First drop)

$$u = 0, h = 90 \text{ m}, g = 10 \text{ m/s}^2$$

by

$$v^2 = u^2 + 2gh$$

$$v^2 = 0 + 2 \times 10 \times 90$$

or

$$v = \sqrt{2 \times 900}$$

$$\text{or } v = 30\sqrt{2} \text{ m/s} = 30 \times 1.414 = 42.420 = 42.4 \text{ m/s}$$

by

$$v = u + gt$$

$$t = \frac{v}{g} = \frac{30\sqrt{2}}{10} = 3\sqrt{2} = 3 \times 1.414 = 4.24 \text{ sec}$$

When ball starts first upward motion -
(First rise)

$$u = 90\% \text{ of } 30\sqrt{2}, v = 0$$

$$= \frac{90}{100} \times 30\sqrt{2} = 27\sqrt{2} = 27 \times 1.414$$

$$\text{or } u = 38.1 \text{ m/s}$$

$$\text{by } v = u + gt$$


$$0 = 38.1 - 10t$$

$$[g \rightarrow -g]$$

$$\text{or } t = 3.81 \text{ sec}$$

First drop ($u=0$)First upward ($v=0$)Second drop ($u=0$)

90 m



When the ball falls second time
(Second drop)

$$t = 3.81 \text{ sec}$$

$$u = 90\% \text{ of } 27\sqrt{2} \text{ (velocity after collision)}$$


$$= \frac{90}{100} \times 27\sqrt{2} = 24.3\sqrt{2} \text{ m/s} = 34.3 \text{ m/s}$$

Now the total time taken by the ball

$$t_T = 4.24 + 3.81 + 3.81$$

$$t_T = 11.86 \text{ sec.}$$

Thereafter body starts second upward motion



$$4.24 + 3.81 = 8.05 \text{ sec}$$

2.9


Topic.....

Date.....

~~3.13~~

(a) Magnitude of displacement can be equal or less than path length but cannot be greater than path length.

e.g.



Suppose a body moves from 'O' to A then 'A' to 'B'. Here displacement = -4 m

and path length = 6 + 6 + 4 = 16 m

We can see mag. of displacement = 4 m

and path length = 16 m

both can be equal for 1d motion only.

(b) For above case let the total time = 10 sec,
Mag. of Average velocity = $\frac{4}{10} = 0.4 \text{ m/s}$

and average speed = $\frac{16}{10} = 1.6 \text{ m/s}$

i.e. average speed can be greater or equal to the mag. of average velocity but cannot be less than mag. of velocity.

both can be equal for 1d motion only.

2.10

~~3.14~~

(a) Given distance of markets = 2.5 km

Speed of man while going to market $V_1 = 5 \text{ km/h}$

" " " " returning to home $V_2 = 7.5 \text{ km/h}$

Time taken to reach the market

$$t_1 = \frac{S}{V_1} = \frac{2.5}{5} = \frac{1}{2} \text{ hr} = 30 \text{ min}$$

Time taken to reach home from market

$$t_2 = \frac{S}{V_2} = \frac{2.5}{7.5} = \frac{1}{3} \text{ hr} = 20 \text{ min}$$

Teacher's Sign

$$\text{Average velocity} = \frac{\text{Total displacement}}{\text{Total time}}$$

and

$$\text{Average Speed} = \frac{\text{Total distance}}{\text{Total time}}$$

(i) 0 to 30 min

In 30 min. man reaches to market.

therefor

$$|\text{Average velocity}| = \frac{2.5}{0.5} = 5 \text{ Km/h}$$

$$\text{Average speed} = \frac{2.5}{0.5} = 5 \text{ Km/h}$$

(ii) 0 to 50 min

In 50 min, man reaches to home.

i.e displacement = 0

So

$$\text{Average velocity} = 0$$

and

$$\text{Average speed} = \frac{2.5 + 2.5}{50/60} \quad [\because 50 \text{ min} = \frac{50}{60} \text{ hr}]$$


$$= \frac{5 \times 6}{5} = 6 \text{ Km/h}$$

(iii) 0 to 40 min

In first 30 min. man reaches market.

In remaining 10 min he covers distance with speed of 7.5 Km/h.

$$\text{i.e } S = 7.5 \times \frac{10}{60} = \frac{7.5}{6} = 1.25 \text{ Km}$$



$$\text{Average velocity} = \frac{\text{Total displacement}}{\text{Total time}}$$

and

$$\text{Average Speed} = \frac{\text{Total distance}}{\text{Total time}}$$

(i) 0 to 30 min

In 30 min. man reaches to market.

therefor

$$|\text{Average velocity}| = \frac{2.5}{0.5} = 5 \text{ km/h}$$

$$\text{Average speed} = \frac{2.5}{0.5} = 5 \text{ km/h}$$

(ii) 0 to 50 min

In 50 min. man reaches to home.

i.e displacement = 0

So

$$\text{Average velocity} = 0$$

and

$$\text{Average speed} = \frac{2.5 + 2.5}{50/60} \quad \left[\because 50 \text{ min} = \frac{50}{60} \text{ hr} \right]$$


$$= \frac{5 \times 6}{5} = 6 \text{ km/h}$$

(iii) 0 to 40 min

In first 30 min. man reaches market.

In remaining 10 min he covers distance with speed of 7.5 km/h.

$$\text{i.e } S = 7.5 \times \frac{10}{60} = \frac{7.5}{6} = 1.25 \text{ km}$$



So |Average velocity| = $\frac{2.5 - 1.25}{40/60}$

$$= \frac{1.25 \times 6^3}{42}$$

$$= \frac{3.75}{2} = 1.875 \text{ km/h}$$

and

Average speed = $\frac{2.5 + 1.25}{40/60}$

$$= \frac{3.75 \times 6^3}{42} = \frac{11.25}{2}$$

$$= 5.625 \text{ km/h}$$

2.11
3.15

Instantaneous velocity of a particle is velocity at any instant of time. In small interval of time |displacement| = distance

Therefore.


|Instantaneous velocity| = Instantaneous speed

$$\text{Instantaneous velocity/speed} = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}$$



2.12
3.16



(a)




(b)



- (a) Not possible because graph shows two different values of position (x) for one value of time.
- (b) Not possible because for one value of time there are two different velocities.
- (c) Not possible because speed cannot be $-ve$.
- (d) Not possible because total path length cannot decrease with time.

2.13
3.17



No, according to graph particle is not in the motion for $t < 0$.

As the graph is for one dimensional motion, it cannot represent parabolic motion. So it is not

correct to say that particle is moving on parabolic path for $t > 0$.

The graph can represent the motion of freely falling particle under gravity.

2.14
3.18

Given,

Speed of police van

$$V_p = 30 \text{ km/h} = \frac{30 \times 5}{18} = \frac{25}{3} \text{ m/s}$$

Speed of thief's car

$$V_t = 192 \text{ km/h} = \frac{192 \times 5}{18} = \frac{160}{3} \text{ m/s}$$

Speed of bullet $V_b = 150 \text{ m/s}$


$\therefore$ Speed of bullet for thief

$$V_{bp} = V_p + V_b$$

($V_b \rightarrow$ speed of bullet)

$$= \frac{25}{3} + 150$$

$$= \frac{475}{3} \text{ m/s}$$



Now relative speed of bullet w.r.to thief's car

$$V_{bt} = V_{bp} - V_t \quad [\because \text{both are in same dir}^n]$$

$$= \frac{475}{3} - \frac{160}{3}$$

$$= \frac{315}{3}$$


$$V_{bt} = 105 \text{ m/s}$$

Hence, bullet hit the thief's car with speed of 105 m/s .

Am

2.15
3.19

(a)



The $x-t$ graph shows that

from t_1 to t_2 particle is at rest.

from t_2 to $t=0$ particle moves with uniform velocity.

from $t=0$ to t_3 particle moves with uniform velocity in opposite direction and reaches its initial position.

from t_3 to t_4 particle continues its uniform motion in -ve x axis.

from t_4 to t_5 particle is in rest again.

Physical situation

The graph may represent a physical situation as a ball at rest on a smooth floor is kicked, it rebounds from a wall with reduced speed and moves to the opposite wall which stops it.

